

**EXTRAPRIN- acetaminophen, aspirin, caffeine tablet**  
**REMEDYREPACK INC.**

*Disclaimer: Most OTC drugs are not reviewed and approved by FDA, however they may be marketed if they comply with applicable regulations and policies. FDA has not evaluated whether this product complies.*

-----

**Extraprin**

***Drug Facts***

***Active ingredients (in each caplet)***

Acetaminophen 250 mg

Aspirin 250 mg (NSAID) \*\*

Caffeine 65 mg

\*\*nonsteroidal anti-inflammatory drug

***Purpose***

Pain reliever

Pain reliever aid

***Uses***

- temporarily relieves minor aches and pains due to:
- headache
- a cold
- arthritis
- muscular aches
- toothache
- premenstrual & menstrual cramps

**Warnings**

**Reye's syndrome:** Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

**Allergy alert :** Acetaminophen may cause severe skin reactions.

Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

**Allergy alert:** Aspirin may cause a severe allergic reaction which may include:

- hives
- facial swelling
- asthma (wheezing)

- shock

**Liver warning:** This product contains acetaminophen. Severe liver damage may occur if you take

- more than 8 caplets in 24 hours, which is the maximum daily amount
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

**Stomach bleeding warning:** This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs [aspirin, ibuprofen, naproxen, or others]
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

**Caffeine warning:** The recommended dose of this product contains about as much caffeine as a cup of coffee. Limit the use of caffeine-containing medications, foods, or beverages while taking this product because too much caffeine may cause nervousness, irritability, sleeplessness, and, occasionally, rapid heartbeat.

#### **Do not use**

- if you have ever had an allergic reaction to acetaminophen, aspirin or other pain reliever/fever reducer
- with any other product containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.

#### **Ask a doctor before use if you have**

- you have liver disease
- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic
- you have asthma

#### **Ask a doctor or pharmacist before use if you are**

- taking a prescription drug for diabetes, gout, or arthritis
- taking any other drug, or are under a doctor's care for any serious condition

#### **Stop use and ask a doctor if**

- an allergic reaction occurs. Seek medical help right away.
- you experience any of the following signs of stomach bleeding:
- feel faint
- vomit blood
- have bloody or black stools
- have stomach pain that does not get better
- ringing in the ears or loss of hearing occurs
- painful area is red or swollen
- pain gets worse or lasts for more than 10 days
- fever gets worse or lasts for more than 3 days
- any new symptoms occur

These could be signs of a serious condition

**If pregnant or breast-feeding,**

ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.

**Keep out of reach of children.**

In case of overdose, get medical help or contact a Poison Control Center (1-800-222-1222) right away. Quick medical attention is critical for adults as well as children even if you do not notice any signs or symptoms.

***Directions***

- **do not use more than directed**
- drink a full glass of water with each dose
- **adults and children 12 years and over:** take 2 caplets every 6 hours; not more than 8 caplets in 24 hours
- **children under 12 years:** ask a doctor

***Other information***

- store at 20°-25°C (68°-77°F)
- close cap tightly after use

***Inactive ingredients***

corn starch, hypromellose, microcrystalline cellulose, povidone, propylene glycol, silicon dioxide, sodium starch glycolate, stearic acid, titanium dioxide, triacetin.

***Questions?***

call toll-free 1-800-935-6737

DRUG: Extraprin

GENERIC: ACETAMINOPHEN, ASPIRIN, CAFFEINE

DOSAGE: TABLET

ADMINISTRATION: ORAL

NDC: 70518-2734-0

COLOR: white

SHAPE: CAPSULE

SCORE: No score

SIZE: 18 mm

IMPRINT: PH052

PACKAGING: 30 in 1 BLISTER PACK

ACTIVE INGREDIENT(S):

- CAFFEINE 65mg in 1
- ASPIRIN 250mg in 1

- ACETAMINOPHEN 250mg in 1

INACTIVE INGREDIENT(S):

- STARCH, CORN
- HYPROMELLOSE, UNSPECIFIED
- MICROCRYSTALLINE CELLULOSE
- POVIDONE
- PROPYLENE GLYCOL
- SILICON DIOXIDE
- STEARIC ACID
- TITANIUM DIOXIDE
- TRIACETIN

## Extraprin Extra Strength

Acetaminophen/Aspirin/Caffeine

**250mg/250mg/65 mg**

**Tablet**

**QTY: 30**



NDC #: 70518-2734-00

Expires:

LOT #:

Source NDC: 54257-0858-02



MFG: Magno-Humphries Labs, Tigard, OR 97223

Keep this and all medication out of the reach of children

Directions For Use: See Package Insert

Store at 20-25°C (68-77°F); excursions permitted to 15-30°C (59-86°F) [See USP]

Repackaged by: RemedyRepack Inc., Indiana, PA 15701, 724.465.8762

**NOT FOR RETAIL SALE**

## EXTRAPRIN

acetaminophen, aspirin, caffeine tablet

### Product Information

Product Type	HUMAN OTC DRUG	Item Code (Source)	NDC:70518-2734(NDC:54257-858)
Route of Administration	ORAL		

### Active Ingredient/Active Moiety

Ingredient Name	Basis of Strength	Strength
ACETAMINOPHEN (UNII: 362O9ITL9D) (ACETAMINOPHEN - UNII:362O9ITL9D)	ACETAMINOPHEN	250 mg
ASPIRIN (UNII: R16CO5Y76E) (ASPIRIN - UNII:R16CO5Y76E)	ASPIRIN	250 mg
CAFFEINE (UNII: 3G6A5W338E) (CAFFEINE - UNII:3G6A5W338E)	CAFFEINE	65 mg

### Inactive Ingredients

Ingredient Name	Strength
STARCH, CORN (UNII: O8232NY3SJ)	
HYPROMELLOSE, UNSPECIFIED (UNII: 3NXW29V3WO)	

MICROCRYSTALLINE CELLULOSE (UNII: OP1R32D61U)				
POVIDONE (UNII: FZ989GH94E)				
PROPYLENE GLYCOL (UNII: 6DC9Q167V3)				
SILICON DIOXIDE (UNII: ETJ7Z6XBU4)				
STEARIC ACID (UNII: 4ELV7Z65AP)				
TITANIUM DIOXIDE (UNII: 15FIX9V2JP)				
TRIACETIN (UNII: XHX3C3X673)				
Product Characteristics				
Color	white	Score	no score	
Shape	CAPSULE	Size	18mm	
Flavor		Imprint Code	PH052	
Contains				
Packaging				
#	Item Code	Package Description	Marketing Start Date	Marketing End Date
1	NDC:70518-2734-0	30 in 1 BLISTER PACK; Type 0: Not a Combination Product	05/18/2020	
Marketing Information				
Marketing Category	Application Number or Monograph Citation		Marketing Start Date	Marketing End Date
OTC monograph not final	part343		05/18/2020	

**Labeler** - REMEDYREPACK INC. (829572556)

Revised: 9/2020

REMEDYREPACK INC.